

Milian MA\*. Histiocitosis de células de Langerhans [*Langerhans's cell histiocytosis*], *Med Oral Patol Oral Cir bucal* 2005;10:185.

\* Profesora Asociada de Medicina Bucal. Universidad de Valencia. España

**CASO CLINICO:** Paciente varón de 42 años , fumador. Es remitido por presentar una úlcera en la encía, de un mes de evolución, que le produce molestias. A la exploración intraoral se observa una úlcera de 0,5 cm sobre un torus mandibular en la encía lingual del 3.6 (fig 1-2). Se solicita una OPT (fig 3), una TAC (fig 4) y se realiza una biopsia (fig 5-6. La figura 6 corresponden a S-100).

**CLINICAL CASE:** A 42 year-old male, smoker, was referred for evaluation of an ulcer located in the gingiva, during one month, that produced disturbances. On the intraoral examination we appreciated an ulceration of 0,5 cm, on a mandibular torus, in the lingual gingiva of the left first molar (Fig. 1-2). A panoramic x-Ray (Fig.3), a CT scan (Fig. 4) and a biopsy were taken (Fig. 5-6. Fig. 6 is a S-100 stain).

**Diagnóstico / Diagnosis**

Histiocitosis de células de Langerhans  
*Langerhans's cell histiocytosis*

