

Journal section: Oral Surgery
 Publication Types: Research

doi:10.4317/medoral.21033
<http://dx.doi.org/doi:10.4317/medoral.21033>

Influence of the prosthetic arm length (palatal position) of zygomatic implants upon patient satisfaction

Hilario Pellicer-Chover ¹, Juan Cervera-Ballester ¹, David Peñarrocha-Oltra ², Leticia Bagán ³, María Peñarrocha-Diago ⁴, Miguel Peñarrocha-Diago ⁵

¹ DDS. Professor of the Master in Oral Surgery and Implantology, Department of Stomatology, Valencia University Medical and Dental School, Valencia, Spain

² DDS, PhD. Associate Professor, Department of Stomatology, Valencia University Medical and Dental School, Valencia, Spain

³ DDS. Collaborator in Oral Medicine, Department of Stomatology, Valencia University Medical and Dental School, Valencia, Spain

⁴ MD, PhD, DDS. Full Professor of Oral Surgery, Department of Stomatology, Valencia University Medical and Dental School, Valencia, Spain

⁵ MD, PhD. Chairman of Oral Surgery, Director of the Master in Oral Surgery and Implantology, Department of Stomatology, Valencia University Medical and Dental School, Valencia, Spain

Correspondence:

Universidad de Valencia
 Clínica Odontológica
 Unidad de Cirugía Bucal
 Calle Gascó Oliag, 1
 46021 - Valencia, Spain
miguel.penarrocha@uv.es

Pellicer-Chover H, Cervera-Ballester J, Peñarrocha-Oltra D, Bagán L, Peñarrocha-Diago MA, Peñarrocha-Diago M. Influence of the prosthetic arm length (palatal position) of zygomatic implants upon patient satisfaction. Med Oral Patol Oral Cir Bucal. 2016 May 1;21 (3):e380-4. <http://www.medicinaoral.com/medoralfree01/v21i3/medoralv21i3p380.pdf>

Article Number: 21033 <http://www.medicinaoral.com/>
 © Medicina Oral S. L. C.I.F. B 96689336 - pISSN 1698-4447 - eISSN: 1698-6946
 eMail: medicina@medicinaoral.com

Indexed in:

Science Citation Index Expanded
 Journal Citation Reports
 Index Medicus, MEDLINE, PubMed
 Scopus, Embase and Emcare
 Índice Médico Español

Received: 01/09/2015
 Accepted: 30/11/2015

Abstract

Background: To assess the influence of the prosthetic arm length (palatal position) of zygomatic implants upon patient comfort and stability, speech, functionality and overall satisfaction.

Material and Methods: A retrospective clinical study was made of patients subjected to rehabilitation of atrophic maxilla with complete maxillary implant-supported fixed prostheses involving a minimum of two zygomatic implants (one on each side) in conjunction with premaxillary implants, and with 12 months of follow-up after implant loading. Subjects used a VAS to score general satisfaction, comfort and stability, speech and functionality, and the results were analyzed in relation to the prosthetic arm length of the zygomatic implants 12 months after prosthetic delivery.

Results: Twenty-two patients participated in the study, receiving 22 prostheses anchored on 148 implants (44 were zygomatic and 94 were conventional implants). The mean right and left prosthetic arm length was 5.9 ± 2.4 mm and 6.1 ± 2.7 mm, respectively, with no statistically significant differences between them ($p=0.576$). The mean scores referred to comfort/retention, speech, functionality and overall satisfaction were high - no correlation being found between prosthetic arm length and patient satisfaction ($p=0.815$).

Conclusions: No relationship could be identified between prosthetic arm length (palatal position) and patient satisfaction.

Key words: Zygomatic implants, patient satisfaction, zygomatic prosthesis, prosthetic arm length.

Introduction

Zygomatic implants, when positioned in conjunction with premaxillary implants, can facilitate the surgical rehabilitation of patients with severe maxillary resorption (1). Furthermore, the final position of the zygomatic implant can affect the configuration of the prosthesis, because of the location and emergence of these implants at a more medial position compared to standard maxillary implants (2).

Some authors (3-6) have reported satisfactory results in patients treated with fixed dental prostheses supported by multiple zygomatic implants. However, Bothur *et al.* (7) observed mild speech deterioration among these patients. Petrovic *et al.* (8) showed that a 2 mm sagittal alteration of the maxillary incisor can cause up to 80% distortion of speech in patients wearing complete dentures, and found that speech distortion increases rapidly when the palatal plate is more than 1 mm thick. The palatal emergence of zygomatic implants placed using the intrasinus technique requires the preparation of a prosthesis with “palatal arms” that invade the lingual space. We have found no studies in the literature on the relationship between the prosthetic arm length (palatal position) of zygomatic implants and patient satisfaction.

The aim of this study was to assess the influence of the prosthetic arm length (palatal position) of zygomatic implants upon patient comfort and stability, speech alterations, functionality, and overall satisfaction.

Material and Methods

The present study is reported in accordance with the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement (9).

- Patient selection

A retrospective clinical study was made in the Oral Surgery and Implant Dentistry Division of the University of Valencia (Valencia, Spain), involving patients treated between 1998 and 2004. The study was carried out following the principles of the Declaration of Helsinki on human research. Accordingly, all patients were informed about the study and were asked to sign an informed consent document before being included. The study design was approved by the ethical review board of the University of Valencia (Ref.: H1402993407096). Patients subjected to rehabilitation of atrophic maxillas (Cawood and Howell Class VI) with complete maxillary implant-supported fixed prostheses involving a minimum of two zygomatic implants (one on each side) in conjunction with premaxillary implants and followed-up on for 12 months after implant loading were included in the study. We excluded patients who failed to complete the questionnaire or who did not attend the follow-up examinations.

- Description of procedures

All surgeries were carried out by the same surgeon

(MPD) under local anesthesia (4% articaine with 1:100,000 adrenalin [Inibsa, Lliça de Vall, Barcelona, Spain]) and sedation (1% propofol solution), with blood pressure, pulse, and oximetric monitoring by the anesthetist. Patients received zygomatic fixtures combined with conventional implants. Zygomatic implants (Nobel Biocare, Göteborg, Sweden) were placed in the malar zygoma using the procedure described by Stella and Warner (2). A further three to 6 conventional implants were inserted into the residual maxillary alveolar process. All implants remained submerged, and the second surgery was carried out three months later.

For the first week following implant placement, patients were instructed not to use provisional prostheses; thereafter they used their pre-existing complete dentures as provisional prostheses during the implant healing period. The definitive screw-retained prostheses were placed four months after surgery.

- Patient-based measurements

All patients were reviewed one month after implant placement and at 6 and 12 months after delivery of the definitive prostheses. The measurements were performed by two independent observers (JCB and HPC) previously calibrated using sample instruments. The concordance study was based on calculation of the weighted kappa index.

We used a line on the occlusal surface of the molars and premolars and cingulum of the maxillary canines and incisors (occlusal line). Then, a perpendicular line was traced running parallel to the palatal prosthetic arm; the line was measured with precision calipers (resolution: 0.01 mm/0.0005” - precision: ± 0.1 mm) (PCE Iberica, S.L. Albacete, Spain) (Fig. 1).

To estimate patient satisfaction with the fixed prosthesis, 10-cm visual analog scales (VAS) (range 1-10) were given to each patient 12 months after prosthesis delivery. The VAS were used to score general satisfaction with the implant-retained prosthesis, comfort and stability, speech, functionality and overall satisfaction. Patients



Fig. 1. View of the reference lines and measurements used in the study of the prostheses.

were asked to rate these aspects of their care on the VAS from 0 (totally dissatisfied) to 10 (completely satisfied). The subjects were asked to draw a vertical line at the point on the horizontal line that best represented their response (4). The patients marked the scale independently in the presence of a research assistant who offered explanations or help as needed.

Results

Twenty-two patients (11 women and 11 men) with a mean age of 54 years (range 31-77) were included in the study. Ten patients presented class IV atrophy and 12 patients had class V atrophy according to the classification of Cawood and Howell. The opposing arch exhibited natural dentition in 5 patients and implant-supported fixed dentures in 5. Five patients had a combination of natural dentition and implant-supported fixed prostheses, and 7 patients already had an overdenture in the opposing arch. Twelve patients received four standard anterior implants, 5 received 6, one received two, three received three, and one patient received 5 implants in the residual maxillary alveolar process (Table 1).

The zygomatic implant success rate was 97.7% (one implant failed during osseointegration and was replaced by a pterygoid implant) and the standard anterior implant success rate was 97.8% (two standard implants failed during the study period).

The mean length of the prosthetic arms was 6 ±2.4 mm (5.9 ±2.4 mm on the right side and 6.1 ±2.7 mm on the left). The overall satisfaction score was 9.45 out of 10 (9.68 for comfort-stability, 9.36 for speech, and 9.64 for functionality).

Since the sample size was small (n=22), normality for the arm length data was checked using the Shapiro-Wilk test (*p*>0.05). Applying a t-test for paired samples, it was confirmed that the mean right side lateral length was equal to that of the left side (*p*=0.576), showing that the longest prosthetic arm and average length were very similar and led to the same results. We decided to base the data analysis on the maximum length (Table 2).

Given that patient satisfaction generated values that were markedly asymmetrical (all patients gave scores of 7 or higher, and most gave scores of 9-10), any possible correlation to length was evaluated using Spearman’s rank correlation coefficient. The results are shown in table 2.

Discussion

In the present study, only two of the 94 anterior implants were lost. The losses occurred during the study period. No other implant failures occurred during the follow-up period, and all prostheses remained in full function. These findings indicate a 97.7% survival rate for the zygomatic implants and a 97.8% survival rate

Table 1. Characteristics of patients treated with zygomatic implants.

Patient	Age	Gender	Smoking *	Atrophy	Opposing dentition	Number of conventional implants
1	58	M	Yes	V	OD	4
2	59	M	No	IV	IFP	6
3	44	F	No	V	OD	4
4	55	F	Yes	V	ND	3
5	68	F	No	IV	ND + IFP	4
6	54	M	No	IV	OD	2
7	54	M	No	IV	IFP	4
8	58	M	Yes	IV	ND	4
9	50	M	Yes	V	ND	4
10	31	M	No	V	ND + IFP	3
11	49	M	No	IV	OD	4
12	46	M	Yes	V	OD	4
13	61	M	Yes	IV	IFP	6
14	54	F	Yes	V	ND + IFP	5
15	54	F	No	IV	IFP	4
16	56	F	No	V	ND	3
17	55	F	Yes	V	ND + IFP	4
18	77	F	No	V	OD	4
19	48	F	No	IV	IFP	6
20	49	M	Yes	IV	ND	6
21	56	F	No	V	ND + IFP	4
22	52	F	No	V	OD	6

Overdenture (OD), implant-supported fixed prosthesis (IFP), natural dentition (ND), natural dentition and implant-supported fixed prosthesis (ND + IFP)

* More than 10 cigarettes per day.

Table 2. Correlations.

			Mean distance	Maximum distance
Spearman's rho	Mean distance	Correlation coefficient	1.000	0.986*
		Significance	-	0.000
	Maximum distance	Correlation coefficient	0.986*	1.000
		Significance	0.671	0.772
	Comfort and stability	Correlation coefficient	-	0.065
		Significance	-	0.772
	Speech	Correlation coefficient	-	0.005
		Significance	-	0.982
	Functionality	Correlation coefficient	-	0.045
		Significance	-	0.843
	Overall	Correlation coefficient	-	0.843
		Significance	-	0.843

Output verifies that mean distance remained identical to data for maximum distance.

*Correlation significance level = 0.01 (two-sided).

for the anterior implants. Previously reported survival rates have ranged from 94.2-100% for zygomatic implants (1,10-12) and from 73-91.8% for anterior implants (1,13,14).

Only two studies (12,14) have evaluated the mean distance from zygomatic head center to ridge center. Aparicio *et al.* (12) obtained a distance on the right and left sides of 4.5 mm and 5.7 mm, respectively. These values were slightly lower than in the present study (5.9 ±2.4 mm on the right side and 6.1 ±2.7 mm on the left side), possible because of the reference points selected for measuring the prosthetic arms.

The present study found no significant differences in patient satisfaction; all patients gave very favorable VAS scores for the implant-supported prostheses, regardless of prosthetic arm length. These results support the findings of other authors (3,4,6,11). However, Aparicio *et al.* (12) observed that four zygomatic abutments in two patients were disconnected because of uncomfortable prostheses, although 84% of patients reported satisfaction levels of over 80%, and 31.8% awarded the maximum score for satisfaction (100%).

Bothur *et al.* (7) observed that during the evaluation period, all patients affirmed their dependence on speech at work, in their spare time, and in daily conversation. Speech is influenced by a number of factors, and difficulties can be initiated by alterations in the length, position, inclination and thickness of the teeth (7). Speech distortion increases rapidly when the palatal plate is made more than 1 mm thicker (8), and the palatal alterations in the present sample were obvious: satisfaction with speech was the lowest of all the parameters

studied, although scores were nevertheless high (9.36 on the VAS). Lundqvist *et al.* (15) observed that following initial phonetic problems, 94% of the individuals considered themselves free from speech problems after three years of follow-up.

Further studies are needed to investigate the position of the implants at palatal level in relation to patient satisfaction with implant-supported fixed dental prostheses.

Conclusions

No relationship was observed between prosthetic arm length and patient satisfaction.

References

1. Peñarrocha M, García B, Martí E, Boronat A. Rehabilitation of severely atrophic maxillae with fixed implant-supported prostheses using zygomatic implants placed using the sinus slot technique: clinical report on a series of 21 patients. *Int J Oral Maxillofac Implants.* 2007;22:645-50.
2. Stella JP, Warner MR. Sinus slot technique for simplification and improved orientation of zygomatic dental implants: a technical note. *Int J Oral Maxillofac Implants.* 2000;15:889-93.
3. Farzad P, Andersson L, Gunnarsson S, Johansson B. Rehabilitation of severely resorbed maxillae with zygomatic implants: an evaluation of implant stability, tissue conditions, and patients' opinion before and after treatment. *Int J Oral Maxillofac Implants.* 2006;21:399-404.
4. Peñarrocha M, Carrillo C, Boronat A, Martí E. Level of satisfaction in patients with maxillary full-arch fixed prostheses: zygomatic versus conventional implants. *Int J Oral Maxillofac Implants.* 2007;22:769-73.
5. Miglioranza RM, Sotto-Maior BS, Senna PM, Francischone CE, Del BelCurry AA. Immediate occlusal loading of extrasinuszygomatic implants: a prospective cohort study with a follow-up period of 8 years. *Int J Oral Maxillofac Surg.* 2012;41:1072-6.

6. Sartori EM, Padovan LE, de MattiasSartori IA, Ribeiro PD Jr, Gomes de Souza Carvalho AC, Goiato MC. Evaluation of satisfaction of patients rehabilitated with zygomatic fixtures. *J Oral Maxillofac Surg.* 2012;70:314-9.
7. Bothur S, Garsten M. Initial speech problems in patients treated with multiple zygomatic implants. *Int J Oral Maxillofac Implants.* 2010;25:379-84.
8. Petrović A. Speech sound distortions caused by changes in complete denture morphology. *J Oral Rehabil.* 1985;12:69-79.
9. Vandembroucke JP, von Elm E, Altman DG, Gøtzsche PC, Mulrow CD, Pocock SJ, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *Epidemiology.* 2007;18:805-35.
10. Bedrossian E, Stumpel L, Beckely M, Indresano T, Indersano T. The zygomatic implant: Preliminary data on treatment of severely resorbed maxillae. A clinical report. *Int J Oral Maxillofac Implants.* 2002;17:861-5.
11. Hirsch JM, Ohnell LO, Henry PJ, Andreasson L, Brånemark PI, Chiapasco M, et al. A clinical evaluation of the Zygoma fixture: one year of follow-up at 16 clinics. *J Oral Maxillofac Surg.* 2004;62(Suppl 2):22-9.
12. Aparicio C, Manresa C, Francisco K, Ouazzani W, Claros P, Potau JM, et al. The long-term use of zygomatic implants: a 10-year clinical and radiographic report. *Clin Implant Dent Relat Res.* 2014;16:447-59.
13. Brånemark PI, Gröndahl K, Ohnell LO, Nilsson P, Petruson B, Svensson B, et al. Zygoma fixture in the management of advanced atrophy of the maxilla: technique and long-term results. *Scand J Plast Reconstr Surg Hand Surg.* 2004;38:70-85.
14. Becktor JP, Isaksson S, Abrahamsson P, Sennerby L. Evaluation of 31 zygomatic implants and 74 regular dental implants used in 16 patients for prosthetic reconstruction of the atrophic maxilla with cross-arch fixed bridges. *Clin Implant Dent Relat Res.* 2005;7:159-65.
15. Lundqvist S, Haraldson T, Lindblad P. Speech in connection with maxillary fixed prostheses on osseointegrated implants: a three-year follow-up study. *Clin Oral Implants Res.* 1992;3:176-80.

Conflict of interest

The authors have declared that no conflict of interest exist.